

A discrete linearizability test based on multiscale analysis

R. Hernández Heredero

Departamento de Matemática Aplicada
Universidad Politécnica de Madrid,
Escuela Universitaria de Ingeniería Técnica de Telecomunicación,
Campus Sur Ctra. de Valencia Km. 7, 28031 Madrid, Spain
E-mail: rafahh@euitt.upm.es

D. Levi

Dipartimento di Ingegneria Elettronica,
Università degli Studi Roma Tre and Sezione INFN, Roma Tre,
Via della Vasca Navale 84, 00146 Roma, Italy
E-mail: levi@roma3.infn.it

C. Scimiterna

Dipartimento di Ingegneria Elettronica,
Università degli Studi Roma Tre and Sezione INFN, Roma Tre,
Via della Vasca Navale 84, 00146 Roma, Italy
E-mail: scimiterna@fis.uniroma3.it

November 2, 2010

Abstract

In this paper we consider the classification of dispersive linearizable partial difference equations defined on a quad-graph by the multiple scale reduction around their harmonic solution. We show that the A_1 , A_2 and A_3 linearizability conditions restrain the number of the parameters which enter into the equation. A subclass of the equations which pass the A_3 C -integrability conditions can be linearized by a Möbius transformation.

1 Introduction

Calogero in 1991 [2] introduced the notion of S and C integrable equations to characterize those nonlinear partial differential equations which are solvable through an Inverse Scattering Transform or linearizable through a change of variables. Using the multiscale reductive technique he was able to show that the Non Linear Schrödinger Equation (*NLSE*)

$$i\partial_t u = K_2[u] = \partial_{xx} u + \rho_2 |u|^2 u, \quad u = u(x, t), \quad (1)$$

appears as a universal equation governing the evolution of slowly varying packets of quasi-monochromatic waves in weakly nonlinear media featuring dispersion. The necessary conditions

for the S -integrability is that ρ_2 is real. If, however, the equation is linearizable then ρ_2 must be null, the equation has to be linear or linearizable, as the Eckhaus equation [4, 20].

By going to a higher order in the expansion, the multiscale techniques have been used to find new S -integrable Partial Differential Equations (PDE's) and to prove the integrability of new nonlinear equations [7, 8, 13]. Probably the most important example of such nonlinear PDE is the Degasperis-Procesi equation [6]. The application of this test to the case of linearizable equations has not been done yet. However Calogero and collaborators constructed many interesting linearizable nonlinear evolution equations by applying complicated transformations to linear equations [3, 5].

In the case of discrete equations it has been shown [1, 9–11, 16–19, 24] that a similar situation is also true. One can present the equivalent of the Calogero-Eckhaus theorem stating that a nonlinear dispersive partial difference equation will not be S -integrable if its multiscale expansion on analytic functions will not give rise to an integrable $NLSE$ at the lowest order. Moreover it was shown on examples that a nonlinear partial difference equation will be C -integrable if its multiscale expansion on analytic functions will give rise to a linear partial differential equation [23].

As was shown in [10, 11, 17], the introduction of multiple scales on a lattice reduces the given discrete equation either to a local nonlinear partial difference equation, by imposing a slow-varying condition, or to a partial differential equation of infinite order when dealing with analytic functions. Here we choose the second alternative because, as was shown in [10, 11], only in this case the integrability conditions of the discrete equation are preserved.

In this paper we provide necessary conditions for the linearizability of a class of real difference equations in the variable $u : \mathbb{Z}^2 \rightarrow \mathbb{R}$ and its three nearest-neighbors defined on a $\mathbb{Z}^2$ square-lattice

$$\mathcal{Q}(u_{n,m}, u_{n+1,m}, u_{n,m+1}, u_{n+1,m+1}; \beta_1, \beta_2, \dots) = 0, \quad (2)$$

where β_i 's are real parameters. Linearizability conditions will be determined through the multiple scale perturbative development.




Figure 1: The $\mathbb{Z}^2$ square-lattice where the equation $\mathcal{Q} = 0$ is defined

We will suppose that eq. (2) is linear-affine in every variable, and our classification will be carried out up to the simultaneous Möbius transformations

$$u_{n,m} \mapsto (Au_{n,m} + B)/(Cu_{n,m} + D). \quad (3)$$

In Section 2 we will briefly discuss the multiple scale expansion for eq. (2) and present the integrability conditions which ensure that the given equation is a C -integrable equation of j order, with $j = 1, 2, 3$, i.e. it is such that asymptotically satisfies the C -integrability conditions up to third order. In Section 3 we apply them to the eq. (2) and present a sequence of theorems which give the conditions on the constants β_i 's under which the system is A_1 , A_2 or A_3 integrable. At the end we present some conclusive remarks.

2 Expansion of real dispersive partial difference equations

We now briefly illustrate all the ingredients of the reductive perturbative technique necessary to treat difference equations, as presented in [9, 10]. We will consider here the multiscale expansion in the case of analytic functions so as to preserve the linearizability of the given equation.

Introducing multiple lattice scales, under some obvious hypotheses on the $\mathcal{C}^{(\infty)}$ property of the function $u_{n,m}$ and on the radius of convergence of its Taylor expansion for all the n and m shifts involved in the difference equation (2), we can write a series representation of the shifted values of $u_{n,m}$ around the point (n, m) . Choosing

$$n_i = \varepsilon_{n_i} n, \quad \varepsilon_{n_i} \doteq N_i \varepsilon^i, \quad 1 \leq i \leq K_n, \quad m_j = \varepsilon_{m_j} m, \quad \varepsilon_{m_j} \doteq M_j \varepsilon^j, \quad 1 \leq j \leq K_m,$$

where the various constants N_i , M_j and ε are all real numbers and we will assume $K_n = 1$ and $K_m = K$ (eventually $K = +\infty$), the total shift operators T_n , T_m can be rewritten in terms of partial shift operators $\mathcal{T}_n$, $\mathcal{T}_m$ as

$$T_n = \mathcal{T}_n \mathcal{T}_{n_1}^{(\varepsilon_{n_1})} = \mathcal{T}_n \sum_{j=0}^{+\infty} \varepsilon^j \mathcal{A}_n^{(j)}, \quad \mathcal{A}_n^{(j)} \doteq \frac{N_1^j}{j!} \partial_{n_1}^j, \quad (4a)$$

$$T_m = \mathcal{T}_m \prod_{j=1}^K \mathcal{T}_{m_j}^{(\varepsilon_{m_j})} = \mathcal{T}_m \sum_{j=0}^{+\infty} \varepsilon^j \mathcal{A}_m^{(j)}, \quad (4b)$$

$$T_n T_m = \mathcal{T}_n \mathcal{T}_m \mathcal{T}_{n_1}^{(\varepsilon_{n_1})} \prod_{j=1}^K \mathcal{T}_{m_j}^{(\varepsilon_{m_j})} = \mathcal{T}_n \mathcal{T}_m \sum_{j=0}^{+\infty} \varepsilon^j \mathcal{A}_{n,m}^{(j)}, \quad (4c)$$

and in terms of differential operators $\mathcal{A}_m^{(j)}$, $\mathcal{A}_{n,m}^{(j)}$. These operators are given by appropriate combinations of $\frac{M_k^j}{j!} \partial_{m_k}^j$ (explicit expressions and formulae can be found in [9], see [22] for more details). We can assume for the function $u_{n,m} = u(n, m, n_1, \{m_j\}_{j=1}^K, \varepsilon)$ a double expansion in harmonics and in the perturbative parameter ε

$$u_{n,m} = \sum_{\gamma=1}^{+\infty} \sum_{\theta=-\gamma}^{\gamma} \varepsilon^\gamma u_\gamma^{(\theta)}(n_1, m_j, j \geq 1) e^{i\theta(\kappa h n - \omega \sigma m)}, \quad (5)$$

with $u_\gamma^{(-\theta)}(n_1, m_j, j \geq 1) = \bar{u}_\gamma^{(\theta)}(n_1, m_j, j \geq 1)$ in order to ensure the reality of $u_{n,m}$. Then, inserting the explicit expressions (4) of the shift operators in terms of the derivatives with respect to the slow variables, eq. (2) turns out to be a PDE of infinite order. Moreover we require the functions $u_\gamma^{(\theta)}$ to satisfy the asymptotic conditions $\lim_{n_1 \rightarrow \pm\infty} u_\gamma^{(\theta)} = 0$, $\forall \gamma$ and θ , and the index γ chosen ≥ 1 in order to let any nonlinear part of eq. (2) to enter as a perturbation in the multiscale expansion.

2.1 The orders beyond the Schrödinger equation and the C-integrability conditions

The multiple scale expansion of a nonlinear partial difference equation on analytic functions will thus give rise to continuous PDE's. So a multiple scale linearizability test will require that an equation is C -integrable if its multiple scale expansion will go into the hierarchy of the Schrödinger equation. To show this we need to consider the orders beyond that at which one obtains for the harmonic $u_1^{(1)}$ the Schrödinger equation. References [7, 14, 15] contain different approaches to the higher order multiscale expansion of S -integrable nonlinear PDE's.

Following [7] we can remove all secular terms from the reduced equations order by order and thus, in agreement with Degasperis and Procesi [8], we can state the following theorem proved by Michela Procesi [21]:

Theorem 1. *A nonlinear dispersive partial difference equation is C -integrable only if its multiscale expansion is given by a uniform asymptotic series such that*

1. *The amplitude $u_1^{(1)}$ evolves at the slow-times m_σ , $\sigma \geq 2$ according to the σ -th equation of the linear Schrödinger hierarchy;*

$$\partial_{m_\sigma} u_1^{(1)} = (-i)^{(\sigma-1)} B_\sigma \partial_{n_1}^\sigma u_1^{(1)}, \quad B_\sigma \doteq -\frac{1}{\sigma!} \frac{d^\sigma \omega(\kappa)}{d\kappa^\sigma}, \quad (6)$$

where B_σ are constants.

2. *The amplitudes of the higher perturbations of the first harmonic $u_j^{(1)}$, $j \geq 2$ evolve at the slow-times m_σ , $\sigma \geq 2$ according to certain linear, nonhomogeneous equations when taking into account the proper asymptotic boundary conditions.*

$$\partial_{m_\sigma} u_j^{(1)} - (-i)^{(\sigma-1)} B_\sigma \partial_{n_1}^\sigma u_j^{(1)} \equiv M_\sigma u_j^{(1)} = f_\sigma(j), \quad (7)$$

$\forall j, \sigma \geq 2$, where $B_\sigma \partial_{n_1}^\sigma u_1^{(j)}$ is the σ -th flow in the linear Schrödinger hierarchy (6). All the other $u_j^{(\theta)}$, $\theta \geq 2$ are expressed in terms of differential monomials of $u_\rho^{(1)}$, $\rho \leq j$.

If the asymptotic expansion is, besides a necessary condition for integrability, also a sufficient condition, has been only conjectured in [21]. The proof of the theorem takes into account that for a C -integrable equation by definition we have a linearizing transformation. The next step is to assume that the function satisfying the linearized equation admits a similar perturbative expansion as in (5) and then to perform a multiscale analysis of the linearizing transformation.

In eq. (7) $f_\sigma(j)$ is a nonhomogeneous *nonlinear* forcing term and B_σ , $\sigma \geq 1$, introduced in eq. (6) are complex constants. Eq. (6) represents a hierarchy of *compatible* evolutions for the function $u_1^{(1)}$. It is obvious that the operators M_σ defined in eq. (7) commute among themselves. Thus, once we fix the index $j \geq 2$ in the set of eqs. (7), their compatibilities imply the following conditions

$$M_\sigma f_{\sigma'}(j) = M_{\sigma'} f_\sigma(j), \quad \forall \sigma, \sigma' \geq 2, \quad (8)$$

where, as $f_\sigma(j)$ and $f_{\sigma'}(j)$ are functions of the different perturbations of the fundamental harmonic up to degree $j-1$, the time derivatives ∂_{t_σ} , $\partial_{t_{\sigma'}}$ of those harmonics appearing respectively in M_σ and $M_{\sigma'}$ have to be eliminated using the evolution equations (6, 7) up to the index $j-1$. The commutativity conditions (8) turn out to be an *integrability test*.

To construct the functions $f_\sigma^{(j)}$ we define homogeneous spaces of functions of $u_j^{(1)}$, its n_1 -derivative and its complex conjugate.

Definition 2.1. A differential monomial $\rho[u_j^{(1)}]$, $j \geq 1$ in the functions $u_j^{(1)}$, its complex conjugate and its n_1 -derivatives is a monomial of “gauge” 1 if it possesses the transformation property

$$\rho[\tilde{u}_j^{(1)}] = e^{i\theta} \rho[u_j^{(1)}], \quad \tilde{u}_j^{(1)} \doteq e^{i\theta} u_j^{(1)};$$

Definition 2.2. A finite dimensional vector space $\mathcal{P}_\nu$, $\nu \geq 2$ is the set of all differential polynomials in the functions $u_j^{(1)}$, $j \geq 1$, their complex conjugates and their n_1 -derivatives of order ν in ε and gauge 1 such that

$$\text{order}\left(\partial_{n_1}^\mu u_j^{(1)}\right) = \text{order}\left(\partial_{n_1}^\mu \bar{u}_j^{(1)}\right) = \mu + j, \quad \mu \geq 0;$$

Definition 2.3. $\mathcal{P}_\nu(\mu)$, $\mu \geq 1$ and $\nu \geq 2$ is the subspace of $\mathcal{P}_\nu$ whose elements are differential polynomials in the functions $u_j^{(1)}$ s, their complex conjugates and their n_1 -derivatives of order ν in ε and gauge 1 for $1 \leq j \leq \mu$.

The basis monomials of the spaces $\mathcal{P}_\nu(\mu)$ can be found, for example, in [22].

The functions $f_\sigma(j)$ belong to a well defined finite dimensional space $\mathcal{P}_\nu(\mu)$ and if the relations (8) are satisfied up to the index k , $k \geq 2$, we say that our equation is asymptotically C -integrable of degree k or A_k C -integrable.

2.2 C -integrability conditions for the Schrödinger hierarchy

In this subsection we present the conditions which must be satisfied for the asymptotic C -integrability of order k or the A_k C -integrability conditions with $k = 1, 2, 3$. To simplify the notation, we will use for $u_j^{(1)}$ the concise form $u(j)$.

The A_1 C -integrability condition is given by the absence of the coefficient ρ_2 of the nonlinear term in the $NLSE$.

The A_2 C -integrability condition is obtained by choosing $j = 2$ in the compatibility conditions (8) with $\sigma = 2$ and $\sigma' = 3$:

$$M_2 f_3(j) = M_3 f_2(j), \quad (9)$$

where $M_2 = \partial_{m_2} + iB_2 \partial_{n_1}^2$ and $M_3 = \partial_{m_3} + B_3 \partial_{n_1}^3$.

In this case $f_2(2) \in \mathcal{P}_4(1)$ and $f_3(2) \in \mathcal{P}_5(1)$ with $\dim(\mathcal{P}_4(1)) = 2$ and $\dim(\mathcal{P}_5(1)) = 5$, so that $f_2(2)$ and $f_3(2)$ will be respectively identified by 2 and 5 complex constants

$$f_2(2) \doteq a u_{n_1}(1) |u(1)|^2 + b \bar{u}_{n_1}(1) u(1)^2, \quad (10a)$$

$$f_3(2) \doteq \alpha |u(1)|^4 u(1) + \beta |u_{n_1}(1)|^2 u(1) + \gamma u_{n_1}(1)^2 \bar{u}(1) + \delta \bar{u}_{n_1 n_1}(1) u(1)^2 + \epsilon |u(1)|^2 u_{n_1 n_1}(1). \quad (10b)$$

In this way, eliminating from eq. (9) the derivatives of $u(1)$ with respect to the slow-times m_2 and m_3 using the evolutions (6) with $\sigma = 2, 3$ and equating term by term, we obtain that the A_2 C -integrability conditions give no constraint for the two coefficients a and b . As a consequence one can say that the A_1 C -integrability condition $\rho_2 = 0$ automatically implies A_2 C -integrability, or that it indeed represents an A_2 C -integrability condition. The expression of $\alpha, \beta, \gamma, \delta, \epsilon$ in terms of a and b are:

$$\alpha = 0, \quad \beta = -\frac{3iB_3 b}{B_2}, \quad \gamma = -\frac{3iB_3 a}{2B_2}, \quad \delta = 0, \quad \epsilon = \gamma. \quad (11)$$

where, for convenience, from now on we will write B for B_3 .

The A_3 C -integrability conditions are derived in a similar way setting $j = 3$ in eq. (9). In this case we have that $f_2(3) \in \mathcal{P}_5(2)$ and $f_3(3) \in \mathcal{P}_6(2)$ with $\dim(\mathcal{P}_5(2)) = 12$ and $\dim(\mathcal{P}_6(2)) = 26$, so that $f_2(3)$ and $f_3(3)$ will be respectively identified by 12 and 26 complex constants

$$\begin{aligned} f_2(3) \doteq & \tau_1 |u(1)|^4 u(1) + \tau_2 |u_{n_1}(1)|^2 u(1) + \tau_3 |u(1)|^2 u_{n_1 n_1}(1) + \tau_4 \bar{u}_{n_1 n_1}(1) u(1)^2 \\ & + \tau_5 u_{n_1}(1)^2 \bar{u}(1) + \tau_6 u_{n_1}(2) |u(1)|^2 + \tau_7 \bar{u}_{n_1}(2) u(1)^2 + \tau_8 u(2)^2 \bar{u}(1) + \tau_9 |u(2)|^2 u(1) \\ & + \tau_{10} u(2) u_{n_1}(1) \bar{u}(1) + \tau_{11} u(2) \bar{u}_{n_1}(1) u(1) + \tau_{12} \bar{u}(2) u_{n_1}(1) u(1), \end{aligned} \quad (12a)$$

$$\begin{aligned} f_3(3) \doteq & \gamma_1 |u(1)|^4 u_{n_1}(1) + \gamma_2 |u(1)|^2 u(1)^2 \bar{u}_{n_1}(1) + \gamma_3 |u(1)|^2 u_{n_1 n_1 n_1}(1) \\ & + \gamma_4 u(1)^2 \bar{u}_{n_1 n_1 n_1}(1) + \gamma_5 |u_{n_1}(1)|^2 u_{n_1}(1) + \gamma_6 \bar{u}_{n_1 n_1}(1) u_{n_1}(1) u(1) \\ & + \gamma_7 u_{n_1 n_1}(1) \bar{u}_{n_1}(1) u(1) + \gamma_8 u_{n_1 n_1}(1) u_{n_1}(1) \bar{u}(1) + \gamma_9 |u(1)|^4 u(2) \\ & + \gamma_{10} |u(1)|^2 u(1)^2 \bar{u}(2) + \gamma_{11} \bar{u}_{n_1}(1) u(2)^2 + \gamma_{12} u_{n_1}(1) |u(2)|^2 + \gamma_{13} |u_{n_1}(1)|^2 u(2) \\ & + \gamma_{14} |u(2)|^2 u(2) + \gamma_{15} u_{n_1}(1)^2 \bar{u}(2) + \gamma_{16} |u(1)|^2 u_{n_1 n_1}(2) + \gamma_{17} u(1)^2 \bar{u}_{n_1 n_1}(2) \\ & + \gamma_{18} u(2) \bar{u}_{n_1 n_1}(1) u(1) + \gamma_{19} u(2) u_{n_1 n_1}(1) \bar{u}(1) + \gamma_{20} \bar{u}(2) u_{n_1 n_1}(1) u(1) \\ & + \gamma_{21} u(2) u_{n_1}(2) \bar{u}(1) + \gamma_{22} \bar{u}(2) u_{n_1}(2) u(1) + \gamma_{23} u_{n_1 i}(2) u_{n_1}(1) \bar{u}(1) \\ & + \gamma_{24} u_{n_1}(2) \bar{u}_{n_1}(1) u(1) + \gamma_{25} \bar{u}_{n_1 i}(2) u_{n_1}(1) u(1) + \gamma_{26} \bar{u}_{n_1}(2) u(2) u(1). \end{aligned} \quad (12b)$$

Let us eliminate from eq. (9) with $j = 3$ the derivatives of $u(1)$ with respect to the slow-times m_2 and m_3 using the evolutions (6) respectively with $\sigma = 2, 3$ and the same derivatives of $u(2)$ using the evolutions (7) with $\sigma = 2, 3$. Equating the remaining terms term by term, the A_3 C -integrability conditions turn out to be:

$$\begin{aligned} \tau_1 &= -\frac{i}{4B_2} [b(\tau_{11} - 2\tau_6) + \bar{a}\tau_7], & \bar{b}\tau_7 &= \frac{1}{2} (b - a) (\tau_{11} + \tau_{10} - \tau_6) + \bar{a}\tau_7, \\ a\tau_8 &= b\tau_8 = 0, & a\tau_9 &= b\tau_9 = 0, & \bar{a}\tau_{12} &= a(\tau_{10} - \tau_{11}) + b\tau_6 + \bar{a}\tau_7, \\ & & (\bar{b} - \bar{a})\tau_{12} &= (b - a)\tau_{10}. \end{aligned} \quad (13)$$

Sometimes a and b turn out to be both real. In this case the conditions given in eqs. (13) become:

$$\begin{aligned} R_1 &= \frac{1}{4B_2} [b(I_{11} - 2I_6) + aI_7], & I_1 &= -\frac{1}{4B_2} [b(R_{11} - 2R_6) + aR_7], \\ (b - a)(R_{11} + R_{10} - R_6 - 2R_7) &= 0, & (b - a)(I_{11} + I_{10} - I_6 - 2I_7) &= 0, \\ (b - a)R_8 &= 0, & (b - a)I_8 &= 0, & (b - a)R_9 &= 0, & (b - a)I_9 &= 0, \\ a(R_{12} + R_{11} - R_{10} - R_7) &= bR_6, & a(I_{12} + I_{11} - I_{10} - I_7) &= bI_6, \\ (b - a)(R_{12} - R_{10}) &= 0, & (b - a)(I_{12} - I_{10}) &= 0, \end{aligned} \quad (14)$$

where $\tau_j = R_j + iI_j$ for $j = 1, \dots, 12$. The expressions of the γ_j as functions of the τ_i are:

$$\begin{aligned}
\gamma_1 &= \frac{3B_3}{4B_2^2} (a\tau_6 - 4iB_2\tau_1 + \bar{b}\tau_{12}), \quad \gamma_2 = \frac{3B_3}{4B_2^2} (b\tau_6 + \bar{a}\tau_7), \\
\gamma_3 &= -\frac{3iB_3\tau_3}{2B_2}, \quad \gamma_4 = 0, \quad \gamma_5 = -\frac{3iB_3\tau_2}{2B_2}, \\
\gamma_6 &= -\frac{3iB_3\tau_4}{B_2}, \quad \gamma_7 = \gamma_5, \quad \gamma_8 = \gamma_3 - \frac{3iB_3\tau_5}{B_2}, \quad \gamma_9 = 0, \quad \gamma_{10} = 0, \\
\gamma_{11} &= 0, \quad \gamma_{12} = -\frac{3iB_3\tau_9}{2B_2}, \quad \gamma_{13} = -\frac{3iB_3\tau_{11}}{2B_2}, \quad \gamma_{14} = 0, \quad \gamma_{15} = -\frac{3iB_3\tau_{12}}{2B_2}, \\
\gamma_{16} &= -\frac{3iB_3\tau_6}{2B_2}, \quad \gamma_{17} = \gamma_{18} = 0, \quad \gamma_{19} = -\frac{3iB_3\tau_{10}}{2B_2}, \quad \gamma_{20} = \gamma_{15}, \quad \gamma_{21} = -\frac{3iB_3\tau_8}{B_2}, \\
\gamma_{22} &= \gamma_{12}, \quad \gamma_{23} = \gamma_{16} + \gamma_{19}, \quad \gamma_{24} = \gamma_{13}, \quad \gamma_{25} = -\frac{3iB_3\tau_7}{B_2}, \quad \gamma_{26} = 0.
\end{aligned}$$

The conditions given in eqs. (13, 14) appear to be new. Their importance resides in the fact that a C -integrable equation must satisfy those conditions.

3 Dispersive affine-linear equations on the square lattice

In this Section we derive the necessary conditions for the linearizability of a dispersive and homogeneous affine-linear equation defined on the square lattice and belonging to the class (2). The most general multilinear equation of the class (2) has at most quartic nonlinearity. Let us introduce into its linear part the solution $u_{n,m} = K^n \Omega^m$ where $K = e^{i\kappa}$ and $\Omega = e^{-i\omega(\kappa)}$. We get that this equation is dispersive if homogeneous and written as

$$\begin{aligned}
\mathcal{Q}_\pm &= a_1(u_{n,m} \pm u_{n+1,m+1}) + a_2(u_{n+1,m} \pm u_{n,m+1}) \\
&+ (\alpha_1 - \alpha_2) u_{n,m} u_{n+1,m} + (\alpha_1 + \alpha_2) u_{n,m+1} u_{n+1,m+1} \\
&+ (\beta_1 - \beta_2) u_{n,m} u_{n,m+1} + (\beta_1 + \beta_2) u_{n+1,m} u_{n+1,m+1} \\
&+ \gamma_1 u_{n,m} u_{n+1,m+1} + \gamma_2 u_{n+1,m} u_{n,m+1} \\
&+ (\xi_1 - \xi_3) u_{n,m} u_{n+1,m} u_{n,m+1} + (\xi_1 + \xi_3) u_{n,m} u_{n+1,m} u_{n+1,m+1} \\
&+ (\xi_2 - \xi_4) u_{n+1,m} u_{n,m+1} u_{n+1,m+1} + (\xi_2 + \xi_4) u_{n,m} u_{n,m+1} u_{n+1,m+1} \\
&+ \zeta u_{n,m} u_{n+1,m} u_{n,m+1} u_{n+1,m+1} = 0,
\end{aligned} \tag{15}$$

where $a_1, a_2 \in \mathbb{R} \setminus \{0\}$, $|a_1| \neq |a_2|$, are the coefficients appearing in the linear part while $\alpha_1, \alpha_2, \beta_1, \beta_2, \gamma_1, \gamma_2, \xi_1, \dots, \xi_4, \zeta$ are eleven real parameters to be determined by using the multi-scale procedure described in Section 2. The linear dispersion relation is given by

$$\omega(\kappa) = \arctan \left[\frac{(a_1^2 - a_2^2) \sin \kappa}{(a_1^2 + a_2^2) \cos \kappa + 2a_1 a_2} \right]. \tag{16}$$

Let us look for those transformations which leave the class of equations $\mathcal{Q}_\pm$ invariant. They will give the equivalence conditions for our classification. It is well known that polynomial equations are invariant under simultaneous Möbius transformations $u_{n,m} \mapsto (Au_{n,m} + B)/(Cu_{n,m} + D)$. However, our class of equations is homogeneous with a restriction on the coefficients of the linear part. After a Möbius transformation a constant term

$$a_0 = B^4 \zeta + 2B^3 D (\xi_1 + \xi_2) + B^2 D^2 [\gamma_1 + \gamma_2 + 2(\alpha_1 + \beta_1)] + 2BD^3 (a_1 + a_2) \tag{17}$$




Figure 2: Representation of the quadratic nonlinearities of $\mathcal{Q}_{\pm}$

will appear, and thus, if we do not want to restrict the coefficients of the equation, we have to set $B = 0$ to have $a_0 = 0$. Under the Möbius transformation with $B = 0$ the coefficients of $\mathcal{Q}_{\pm}$ become

$$a_1 \mapsto D^3 a_1, \quad a_2 \mapsto D^3 a_2, \quad \alpha_1 \mapsto D^2 [\alpha_1 + C(a_1 + a_2)], \quad \alpha_2 \mapsto D^2 \alpha_2, \quad (18)$$

$$\beta_1 \mapsto D^2 [\beta_1 + C(a_1 + a_2)], \quad \beta_2 \mapsto D^2 \beta_2, \quad (19)$$

$$\gamma_1 \mapsto D^2 (\gamma_1 + 2C a_1), \quad \gamma_2 \mapsto D^2 (\gamma_2 + 2C a_2), \quad (20)$$

$$\xi_1 \mapsto D\xi_1 + \frac{1}{2}CD [3C(a_1 + a_2) + \gamma_1 + \gamma_2 + 2(\alpha_1 - \alpha_2 + \beta_1)], \quad (21)$$

$$\xi_2 \mapsto D\xi_2 + \frac{1}{2}CD [3C(a_1 + a_2) + \gamma_1 + \gamma_2 + 2(\alpha_1 + \alpha_2 + \beta_1)], \quad (22)$$

$$\xi_3 \mapsto D\xi_3 + \frac{1}{2}CD [C(a_1 - a_2) + \gamma_1 - \gamma_2 + 2\beta_2], \quad (23)$$

$$\xi_4 \mapsto D\xi_4 + \frac{1}{2}CD [C(a_1 - a_2) + \gamma_1 - \gamma_2 - 2\beta_2], \quad (24)$$

$$\zeta \mapsto \zeta + C^2 [2C(a_1 + a_2) + \gamma_1 + \gamma_2 + 2(\alpha_1 + \beta_1)] + 2C(\xi_1 + \xi_2). \quad (25)$$

So our equivalence transformation with respect to which we will be classifying the equation $\mathcal{Q}_{\pm}$ is a restricted Möbius transformation of the form $u_{n,m} \mapsto u_{n,m}/(Cu_{n,m} + D)$.

In this article we limit ourselves to consider the case of $\mathcal{Q}_{+}$. The case $\mathcal{Q}_{-}$ will be dealt with in a subsequent publication.

Imposing that $\rho_2 = 0$ we get the following Proposition:

Proposition 1. *The lowest order necessary conditions for the linearizability of equations $\mathcal{Q}_{+}$ give 6 different classes of equations characterized by different non-superimposed ranges of values of the coefficients of the equation $\mathcal{Q}_{+}$. They are:*

- *Case L1:*

$$\left\{ \begin{array}{l} \alpha_2 = \beta_2 = 0, \quad \alpha_1 = \beta_1, \quad \gamma_1 + \gamma_2 = 2\beta_1, \\ \xi_1 = \xi_2 = \frac{(a_1 + a_2)^2 \gamma_1 \gamma_2 + (3a_1 - 2a_2)a_2 \gamma_1 \beta_1 - a_1(2a_1 - 3a_2)\gamma_2 \beta_1}{4a_1 a_2 (a_1 + a_2)}, \\ \xi_3 = \xi_4 = \frac{-(a_1 - a_2)(a_1 + a_2)^2 \gamma_1 \gamma_2 - a_2(-a_1^2 - 5a_2 a_1 + 2a_2^2)\gamma_1 \beta_1 + a_1(2a_1^2 - 5a_2 a_1 - a_2^2)\gamma_2 \beta_1}{4a_1 a_2 (a_1 + a_2)^2}. \end{array} \right. \quad (26)$$

- *Case L2:*

$$\begin{cases} \alpha_2 = \beta_2 = 0, & \alpha_1 = \beta_1, & (3a_1 - 2a_2)a_2^2\gamma_1 + a_1^2(2a_1 - 3a_2)\gamma_2 = 4a_1(a_1 - a_2)a_2\beta_1, \\ \xi_1 = \xi_2 = \frac{(a_1 + a_2)(a_2^2\gamma_1^2 - a_1^2\gamma_2^2) + 2a_2(a_1^2 - 2a_2^2)\gamma_1\beta_1 + 2a_1(2a_1^2 - a_2^2)\gamma_2\beta_1 - 6a_1(a_1 - a_2)a_2\beta_1^2}{4a_1a_2(a_1^2 - a_2^2)}, \\ \xi_3 = \xi_4 = \frac{2a_1a_2(a_1 + a_2)(\gamma_1 - \gamma_2)\beta_1 + (a_1 - a_2)(a_2\gamma_1 - a_1\gamma_2)^2 + 2a_1a_2(a_2 - a_1)\beta_1^2}{4a_1a_2(a_1 + a_2)^2}. \end{cases} \quad (27)$$

- *Case L3:*

$$\begin{cases} \alpha_2 = \beta_2 = 0, & a_1 = 2a_2, & \gamma_1 = \frac{2}{3}(\alpha_1 + \beta_1), & \gamma_2 = \frac{1}{3}(\alpha_1 + \beta_1), \\ \xi_1 = \xi_2 = -\frac{-5\alpha_1\beta_1 + \alpha_1^2 + \beta_1^2}{6a_2}, & \xi_3 = \xi_4 = -\frac{(\alpha_1 - 2\beta_1)(2\alpha_1 - \beta_1)}{18a_2}. \end{cases} \quad (28)$$

- *Case L4:*

$$\begin{cases} \alpha_2 = \beta_2 = 0, & 2a_1 = a_2, & \gamma_1 = \frac{1}{3}(\alpha_1 + \beta_1), & \gamma_2 = \frac{2}{3}(\alpha_1 + \beta_1), \\ \xi_1 = \xi_2 = -\frac{-5\alpha_1\beta_1 + \alpha_1^2 + \beta_1^2}{6a_1}, & \xi_3 = \xi_4 = \frac{(\alpha_1 - 2\beta_1)(2\alpha_1 - \beta_1)}{18a_1}. \end{cases} \quad (29)$$

- *Case L5:*

$$\begin{cases} \alpha_2 = \beta_2, & \alpha_1 = \beta_1, & 2a_1 = a_2, & \gamma_1 = \frac{2\beta_1}{3}, & \gamma_2 = \frac{4\beta_1}{3}, \\ \xi_1 = \frac{3\beta_1^2 - 2\beta_2\beta_1 + \beta_2^2}{6a_1}, & \xi_2 = \frac{3\beta_1^2 + 2\beta_2\beta_1 + \beta_2^2}{6a_1}, \\ \xi_3 = -\frac{\beta_1^2 - 6\beta_2\beta_1 + 7\beta_2^2}{18a_1}, & \xi_4 = -\frac{\beta_1^2 + 6\beta_2\beta_1 + 7\beta_2^2}{18a_1}. \end{cases} \quad (30)$$

- *Case L6:*

$$\begin{cases} \alpha_2 = -\beta_2, & \alpha_1 = \beta_1, & a_1 = 2a_2, & \gamma_1 = \frac{4\beta_1}{3}, & \gamma_2 = \frac{2\beta_1}{3}, \\ \xi_1 = \frac{3\beta_1^2 + 2\beta_2\beta_1 + \beta_2^2}{6a_2}, & \xi_2 = \frac{3\beta_1^2 - 2\beta_2\beta_1 + \beta_2^2}{6a_2}, \\ \xi_3 = \frac{\beta_1^2 + 6\beta_2\beta_1 + 7\beta_2^2}{18a_2}, & \xi_4 = \frac{\beta_1^2 - 6\beta_2\beta_1 + 7\beta_2^2}{18a_2}. \end{cases} \quad (31)$$

The corresponding six subclasses of equations are invariant under restricted Möbius transformations of the form $u_n \mapsto u_n / (Cu_n + D)$.

Proof: Following the procedure described in Section 2 we expand the fields appearing in equation Q_+ using the definitions (5, 4). The lowest order necessary conditions for the integrability of Q_+ are obtained at the order ε^3 of the multiple scale expansion. At this order we get the m_2 -evolution equation for the harmonic $u_1^{(1)}$, that is in general a nonlinear Shrödinger equation of the form

$$i\partial_{m_2}u_1^{(1)} - B_2\partial_{n_2}^2u_1^{(1)} - \rho_2u_1^{(1)}|u_1^{(1)}|^2 = 0, \quad \text{where,} \quad n_2 = n_1 - \frac{d\omega}{d\kappa}m_1, \quad (32)$$

where the coefficients B_2 and ρ_2 will depend on the parameters of the equation Q_+ and on the wave parameters κ and ω , with ω expressed in terms of κ through the dispersion relation (16).

Let us give just a sketch of the construction of eq. (32), omitting all intermediate formulas. At $\mathcal{O}(\varepsilon)$ we get:

- for $\theta = 1$ a linear equation which is identically satisfied by the dispersion relation (16).
- for $\theta = 0$ a linear equation whose solution is $u_1^{(0)} = 0$.

At $\mathcal{O}(\varepsilon^2)$, taking into account the dispersion relation (16), we get:

- for $\theta = 2$ an algebraic relation between $u_2^{(2)}$ and $u_1^{(1)}$.
- for $\theta = 1$ a linear wave equation for $u_1^{(1)}$, whose solution is given by $u_1^{(1)}(n_1, m_1, m_2) = u_1^{(1)}(n_2, m_2)$, with n_2 given by eq. (32).
- for $\theta = 0$ an algebraic relation between $u_1^{(1)}$ and $u_2^{(0)}$.

Let us stress here that at $\mathcal{O}(\varepsilon^2)$ we find that all the harmonics will depend on the slow-variables n_1 and m_1 through n_2 .

At $\mathcal{O}(\varepsilon^3)$, for $\theta = 1$, by using the previous results, one gets the *NLSE* (32) with

$$B_2 = -\frac{a_1 a_2 (a_1^2 - a_2^2) \sin \kappa}{(a_1^2 + a_2^2 + 2a_1 a_2 \cos \kappa)^2}, \quad \rho_2 = \mathcal{R}_1 + i\mathcal{R}_2,$$

where

$$\mathcal{R}_1 = \frac{\sin \kappa [\mathcal{R}_1^{(0)} + \mathcal{R}_1^{(1)} \cos \kappa + \mathcal{R}_1^{(2)} \cos^2 \kappa + \mathcal{R}_1^{(3)} \cos^3 \kappa + \mathcal{R}_1^{(4)} \cos^4 \kappa]}{(a_1 + a_2)(a_1^2 + a_2^2 + 2a_1 a_2 \cos \kappa)^2 [(a_1 - a_2)^2 + 2a_1 a_2 \cos \kappa (1 + \cos \kappa)]}, \quad (33)$$

$$\mathcal{R}_2 = \frac{\mathcal{R}_2^{(0)} + \mathcal{R}_2^{(1)} \cos \kappa + \mathcal{R}_2^{(2)} \cos^2 \kappa + \mathcal{R}_2^{(3)} \cos^3 \kappa + \mathcal{R}_2^{(4)} \cos^4 \kappa + \mathcal{R}_2^{(5)} \cos^5 \kappa}{(a_1 + a_2)(a_1^2 + a_2^2 + 2a_1 a_2 \cos \kappa)^2 [(a_1 - a_2)^2 + 2a_1 a_2 \cos \kappa (1 + \cos \kappa)]}. \quad (34)$$

Here the coefficients $\mathcal{R}_1^{(i)}$, $0 \leq i \leq 4$, and $\mathcal{R}_2^{(i)}$, $0 \leq i \leq 5$, are polynomials depending on the coefficients $a_1, a_2, \alpha_1, \alpha_2, \beta_1, \beta_2, \gamma_1, \gamma_2, \xi_1, \dots, \xi_4$ and their expressions are cumbersome, so that we omit them.

Note that B_2 is a real coefficient depending only on the parameters of the linear part of $\mathcal{Q}_+$, while ρ_2 is a complex one. Hence the linearizability of the *NLS* equation (32) is equivalent to the request $\rho_2 = 0 \forall \kappa$, that is

$$\mathcal{R}_j^{(i)} = 0, \quad 0 \leq i \leq 5, j = 1, 2. \quad (35)$$

Eq. (35) is a nonlinear algebraic system of eleven equations in twelve unknowns, the coefficient ζ not appearing at this order of the multiple scale expansion. By solving it with the help of the computer algebra software Mathematica one gets the solutions (26–31). These solutions are computed taking into account that $a_1, a_2 \in \mathbb{R} \setminus \{0\}$ with $|a_1| \neq |a_2|$. Let us point out that the equations have been solved using only rational algebra, avoiding the unreliability of computer algebra calculations when using irrational functions.

One first solves the six equations corresponding to $\mathcal{R}_2^{(i)} = 0$, $0 \leq i \leq 5$. Two of them can be solved for ξ_1 and ξ_3 in (rational) terms of the remaining ten coefficients. The resulting system of equations turns out to be ξ_j -independent and linear in the four variables $\alpha_1, \beta_1, \gamma_1$ and γ_2 . Therefore we may write the remaining four equations as a matrix equation with coefficients nonlinearly depending on α_2, β_2, a_1 and a_2 . The rank of the matrix is three. Solutions (26–31) are obtained by requiring that the matrix be of rank 3, 2, 1 and 0. In this way we get

- Case 1:

$$\begin{cases} \alpha_2 = \beta_2 = 0, \\ \xi_1 = \xi_2, \quad \xi_3 = \xi_4. \end{cases} \quad (36)$$

- Case 2:

$$\begin{cases} \alpha_2 = \beta_2, & \alpha_1 = \beta_1, \\ a_1 = 2a_2, \\ \gamma_1 = 2\gamma_2, \\ a_1(\xi_1 - \xi_2) = -a_1(\xi_3 - \xi_4) = -2\alpha_2\gamma_2. \end{cases} \quad (37)$$

- Case 3:

$$\begin{cases} \alpha_2 = -\beta_2, & \alpha_1 = \beta_1, \\ a_2 = 2a_1, \\ \gamma_2 = 2\gamma_1, \\ a_1(\xi_1 - \xi_2) = a_1(\xi_3 - \xi_4) = -\alpha_2\gamma_1. \end{cases} \quad (38)$$

- Case 4:

$$\begin{cases} \alpha_1 = \beta_1 = \frac{1}{2}(1 + a_1/a_2)\gamma_2, \\ a_2\gamma_1 = a_1\gamma_2, \\ a_1(\xi_1 - \xi_2) = -\alpha_2\gamma_1, \\ a_1(\xi_3 - \xi_4) = \beta_2\gamma_1. \end{cases} \quad (39)$$

- Case 5:

$$\begin{cases} (a_2 - a_1)\beta_2 = (a_2 + a_1)\alpha_2, \\ 2a_1a_2(a_1 - a_2)\alpha_1 = (a_1 + a_2)(\gamma_2a_1^2 - \gamma_1a_2^2), \\ 2a_1a_2\beta_1 = \gamma_1a_2^2 + \gamma_2a_1^2, \\ (a_2 - a_1)(\xi_1 - \xi_2) = (\gamma_1 - \gamma_2)\alpha_2, \\ (a_2 - a_1)^2(\xi_3 - \xi_4) = [\gamma_2(a_2 - 3a_1) - \gamma_1(a_1 - 3a_2)]\alpha_2. \end{cases} \quad (40)$$

- Case 6:

$$\begin{cases} (a_2 + a_1)\beta_2 = (a_2 - a_1)\alpha_2, \\ 2a_1a_2\alpha_1 = \gamma_1a_2^2 + \gamma_2a_1^2, \\ 2a_1a_2(a_1 - a_2)\beta_1 = (a_1 + a_2)(\gamma_2a_1^2 - \gamma_1a_2^2), \\ (a_2^2 - a_1^2)(\xi_1 - \xi_2) = [\gamma_1(a_1 - 3a_2) - \gamma_2(a_2 - 3a_1)]\alpha_2, \\ (a_1 + a_2)(\xi_3 - \xi_4) = (\gamma_2 - \gamma_1)\alpha_2. \end{cases} \quad (41)$$

Then we impose the remaining five equations $\mathcal{R}_1^{(i)} = 0$, $0 \leq i \leq 4$ to each of the six obtained solutions (36–41). By a straightforward computer aided computation we conclude that Case 1 has four subcases, Cases (26–29), that pass those conditions while Case 4 has two subcases, (30, 31). All the other cases do not satisfy the equations $\mathcal{R}_1^{(i)} = 0$, $0 \leq i \leq 4$. Cases 1 and 4 represent the necessary and sufficient conditions for the coefficients a and b in (10a) to be real. A direct calculation proves invariance of the resulting equations with respect to the reduced Möbius transformation. Let us stress again that Cases (26–31) are A_2 C -integrable. $\square$

As a consequence of this proposition we can state the following obvious but important Corollary:

Corollary 1. *If the coefficients $a_1, a_2, \alpha_1, \alpha_2, \beta_1, \beta_2, \gamma_1, \gamma_2, \xi_1, \dots, \xi_4$ of equations $\mathcal{Q}_+$ do not satisfy the conditions in Eqs. (26–31) then $\mathcal{Q}_+$ is not linearizable.*

Note that the trivial linearizability condition $\alpha_1 = \alpha_2 = \beta_1 = \beta_2 = \gamma_1 = \gamma_2 = \xi_1 = \xi_2 = \xi_3 = \xi_4 = 0$ is contained in eqs. (26–31) (cases L1 to L6).

A particularly interesting case is when we consider an equation with at most quadratic non-linearity. In this case we get the following Proposition:

Proposition 2. *If $\xi_1 = \xi_2 = \xi_3 = \xi_4 = 0$ in equations $\mathcal{Q}_+$ then the lowest linearizability conditions are:*

- *Cases QL1a and QL1b: $\alpha_2 = \beta_2 = 0$, $\alpha_1 = \beta_1$, $2a_1^2 - 7a_2a_1 + 2a_2^2 = 0 \Leftrightarrow a_1 = \frac{1}{4}(7 \pm \sqrt{33})a_2$, $\gamma_1 = \left(1 \mp \sqrt{\frac{3}{11}}\right)\beta_1$, $\gamma_2 = \left(1 \pm \sqrt{\frac{3}{11}}\right)\beta_1$.*
- *Cases QL2a and QL2b: $\alpha_2 = \beta_2 = 0$, $\alpha_1 = \beta_1$, $4a_1^4 - 12a_2a_1^3 + 9a_2^2a_1^2 - 12a_2^3a_1 + 4a_2^4 = 0 \Leftrightarrow a_1 = \left(\frac{3}{4} + \frac{1}{\sqrt{2}} \pm \frac{1}{2}\sqrt{\frac{1}{4} + 3\sqrt{2}}\right)a_2$, $\gamma_1 = \mp 2\sqrt{\frac{1}{41}(19 + 18\sqrt{2})}\beta_1$, $\gamma_2 = \pm 2\sqrt{\frac{1}{41}(19 + 18\sqrt{2})}\beta_1$.*

Cases QL1a, QL1b are in case L1 and cases QL2a, QL2b are in case L2.

Let us consider now the higher order terms of the expansion for the cases L1–L6. Imposing the A_3 C –integrability conditions (14) on the real and imaginary parts of the coefficients τ_j , $j = 1, \dots, 12$, defined in the expression (12a), one obtains that the most general A_3 C –integrable equation is represented by:

$$\begin{aligned} \alpha_1 = \beta_1 &= \frac{(a_1 + a_2)\gamma_1}{2a_1}, & \alpha_2 = \beta_2 &= 0, & \gamma_2 &= \frac{a_2\gamma_1}{a_1}, \\ \xi_1 = \xi_2 &= \frac{3(a_1 + a_2)\gamma_1^2}{8a_1^2}, & \xi_3 = \xi_4 &= \frac{(a_1 - a_2)\gamma_1^2}{8a_1^2}. \end{aligned} \quad (42)$$

The case (42) is the intersection of the cases (26) and (27). As a consequence of the result (42) we have that an equation of the form $\mathcal{Q}_+$ which satisfies the Proposition 2, i.e. dispersive with at most quadratic nonlinearity, will never be C –integrable.

Moreover it is easy to prove the following theorem:

Theorem 2. *The equations $\mathcal{Q}_+$ satisfying the A_3 C –integrability (42) can be linearized by a real Möbius transformation*

$$u_{n,m} = \frac{\alpha v_{n,m} + \beta}{\gamma v_{n,m} + \delta}$$

if and only if $\zeta = \frac{(a_1 + a_2)\gamma_1^3}{4a_1^3}$. The coefficients of the linearizing transformation are

$$\beta = 0, \quad \gamma = -\frac{\alpha\gamma_1}{2a_1},$$

while the resulting linearized equation is

$$v_{n,m} + v_{n+1,m+1} + \frac{a_2}{a_1}(v_{n+1,m} + v_{n,m+1}) = 0. \quad (43)$$

Equation (43) is the most general linear dispersive equation defined on the square. As the Möbius transformation used is restricted Möbius transformation, (43) can be assumed to be the canonical equation for this case.

4 Conclusions

In this work we presented the complete classification of the linerizable dispersive partial difference equations belonging to the $\mathcal{Q}_+$ class using multiple scales expansions around a periodic

discrete wave solution of the linearized equation, up to the fifth order in the perturbation expansion parameter. The resulting linearizable system depends on three free parameters, and requesting the resulting equation to explicitly linearize via a Möbius transformation reduces the free parameters to one. It is still an open problem the proof of the linearizability of the equation $\mathcal{Q}_+$ satisfying just the A_3 C -integrability conditions (42). Maybe by going higher in the perturbation expansion we could be able to fix the parameter ζ according to the theorem of last Section.

This calculation shows that the multiple scale expansion can be effectively used to classify discrete equations.

Work is in progress for the derivation of the integrable class of dispersive partial difference equations belonging to the $\mathcal{Q}_+$ class. They are provided by a reduced set of equations with respect to the one considered in this case, as $\rho_2 \neq 0$ is just a real constant. Moreover the integrability conditions are given by a larger number of nonlinear algebraic equations for the parameters entering into the equation and, as such, are much harder to solve.

Work is also in progress in the study of the $\mathcal{Q}_-$ case. In this case we get at the lowest order in the perturbative parameter a nonlinear system of partial differential equations relating the zeroth and the fundamental harmonic. The solution of this equation is the key ingredient in the classification of this class of equations which contains all dispersive equations belonging to the ABS classification of multilinear equations on the square.

Acknowledgments

LD and SC have been partly supported by the Italian Ministry of Education and Research, PRIN “Nonlinear waves: integrable fine dimensional reductions and discretizations” from 2007 to 2009 and PRIN “Continuous and discrete nonlinear integrable evolutions: from water waves to symplectic maps” from 2010. RHH thanks the INFN, Sezione Roma Tre and the UPM for their support during his visits to Rome.

References

- [1] M. Agrotis, S. Lafortune and P.G. Kevrekidis, *On a discrete version of the Korteweg-de Vries equation*. Discr. Cont. Dyn. Sist. (2005) suppl. 22–29.
- [2] F. Calogero, *Why are certain nonlinear PDEs both widely applicable and integrable?* in *What is integrability?*, edited by V.E. Zakharov, Springer, Berlin, 1991, 1–62.
- [3] F. Calogero, *Classical Many-Body Problems Amenable to Exact Treatments: (Solvable and/or Integrable and/or Linearizable...) in One-, Two- and Three-Dimensional Space* Springer, New York 2001.
- [4] F. Calogero and W. Eckhaus, *Necessary conditions for integrability of nonlinear PDEs*. Inv. Prob. 3 (1987) L27–L32 ; F. Calogero and W. Eckhaus, *Nonlinear evolution equations, rescalings, model PDEs and their integrability. I*. Inv. Prob. 3 (1987) 229–262; F. Calogero and W. Eckhaus, *Nonlinear evolution equations, rescalings, model PDEs and their integrability. II*. Inv. Prob. 4 (1987) 11–33; F. Calogero, A. Degasperis and X-D. Ji, *Nonlinear Schrödinger-type equations from multiscale reduction of PDEs. II. Necessary conditions of*

- integrability for real PDEs.* Jour. Math. Phys. 42 (2001) 2635–2652; F. Calogero, A. Degasperis and X-D. Ji, *Nonlinear Schrödinger-type equations from multiscale reduction of PDEs. I. Systematic derivation.* Jour. Math. Phys. 41 (2000) 6399–6443; F. Calogero and A. Maccari, *Equations of nonlinear Schrödinger type in $1+1$ and $2+1$ dimensions obtained from integrable PDEs.*, in *Inverse problems: an interdisciplinary study*, Adv. Electron. Electron Phys., Suppl. 19, C.P. Sabatier ed., Academic Press, London, 1987, 463–480.
- [5] F. Calogero and Xiaoda Ji, *C-integrable nonlinear partial differential equations. I* J. Math. Phys. **32** (1991) 87587, *C-integrable nonlinear PDEs. II* J. Math. Phys. **32** (1991) 270317, *C-integrable nonlinear partial differential equations. III* J. Math. Phys. **34** (1993) 581031, *C-integrable nonlinear PDEs. IV* J. Phys. A: Math. Gen. **29** (1996) 67816794.
- [6] A. Degasperis, D.D. Holm, and A.N.I. Hone, *A new integrable equation with peakon solutions.*, (Russian) Teoret. Mat. Fiz. 133 (2002) 170–183; translation in Theoret. and Math. Phys. 133 (2002) 1463–1474.
- [7] A. Degasperis, S.V. Manakov and P.M. Santini, *Multiple-scale perturbation beyond the nonlinear Schrödinger equation. I.*, Phys. D 100 (1997) 187–211.
- [8] A. Degasperis and M. Procesi, *Asymptotic integrability.* in *Symmetry and perturbation theory, SPT98*, edited by A. Degasperis and G. Gaeta, World Scientific, Singapore, 1999, 23–37;
A. Degasperis, *Multiscale expansion and integrability of dispersive wave equations*, in *Integrability*, A.V. Mikhailov ed., Springer, Berlin, 2009, 215–244.
- [9] R. Hernandez Heredero, D. Levi, M. Petrera and C. Scimiterna, *Multiscale expansion of the lattice potential KdV equation on functions of an infinite slow-varyness order*, J. Phys. A, 40 (2007) F831–F840.
- [10] R. Hernandez Heredero, D. Levi, M. Petrera and C. Scimiterna, *Multiscale expansion on the lattice and integrability of partial difference equations*, J. Phys. A, 41 (2008) 315208–315219.
- [11] R. Hernandez Heredero, D. Levi, M. Petrera and C. Scimiterna, *Multiscale expansion and integrability properties of the lattice potential KdV equation*, J. Nonl. Math. Phys., 15 (2008) 313–323.
- [12] T. Kano, “Normal form of nonlinear Schrödinger equation”, J.Phys.Soc.Japan, 58, 4322, 1989.
- [13] Y. Kodama and A.V. Mikhailov, *Obstacles to asymptotic integrability*, in *Algebraic aspects of integrable systems*, Birkhäuser, Boston, MA, 1996, 173–204 and
Y. Hiraoka and Y. Kodama, *Normal Form and Solitons*, in *Integrability*, A.V. Mikhailov ed., Springer, Berlin, 2009, pp. 175–214.
- [14] Y. Kodama, *On integrable systems with higher order corrections*, Phys. Lett. **A 107** (1985) 245–249.
- [15] Y. Kodama, *Nearly integrable systems: normal form and solitons*, in *Nonlinear evolutions*, J. J. P. Leon ed., World Sci. Publ., Teaneck, NJ, 1988, pp. 559–570.

- [16] J. Leon and M. Manna, *Multiscale analysis of discrete nonlinear evolution equations*. Jour. Phys. A 32 (1999) 2845–2869.
- [17] D. Levi, *Multiple-scale analysis of discrete nonlinear partial difference equations: the reduction of the lattice potential KdV*. Jour. Phys. A, 38 (2005) 7677–7689.
- [18] D. Levi and R. Hernandez Heredero, *Multiscale analysis of discrete nonlinear evolution equations: the reduction of the dNLS*. Jour. Nonlinear Math. Phys., 12 suppl. 1(2005) 440–448.
- [19] D. Levi and M. Petrera, *Discrete reductive perturbation technique*. Jour. Math. Phys., 47 (2006) 043509, 20 pp.
- [20] D. Levi and C. Scimiterna, *The Kundu-Eckhaus equation and its discretizations*, J. Phys. A: Math. Theor. **42** (2009) 465203 (8pp).
- [21] M. Procesi, *Non linear waves, multiscale methods and integrability* master degree thesis, Physics Department, “La Sapienza” University, Rome 1997.
- [22] C. Scimiterna, *Multiscale techniques for nonlinear difference equations*, Ph.D. diss., Roma Tre University 2009. <http://dspace-roma3.caspur.it/handle/2307/408f>
- [23] C. Scimiterna and D. Levi, *C-Integrability Test for Discrete Equations via Multiple Scale Expansions*, SIGMA **6** (2010) 070 17 pages.
- [24] S.W. Schoombie, *A discrete multiple scales analysis of a discrete version of the Korteweg-de Vries equation*. Jour. Comp. Phys., 101 (1992) 55–70.